

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C07K 7/56, A61K 38/12</b>		A1	(11) International Publication Number: <b>WO 97/17365</b>
			(43) International Publication Date: 15 May 1997 (15.05.97)
(21) International Application Number: PCT/US96/17715		(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 6 November 1996 (06.11.96)			
(30) Priority Data: 60/006,505 9 November 1995 (09.11.95) US 9603241.2 16 February 1996 (16.02.96) GB			
(71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).		<p><b>Published</b></p> <p><i>With international search report.</i></p> <p><i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
(72) Inventors; and			
(75) Inventors/Applicants (for US only): BALKOVEC, James, M. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). BOUFFARD, Frances, A. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). DROPINSKI, James, F. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).			
(74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).			

(54) Title: CYCLOHEXAPEPTIDYL BISAMINE COMPOUND, COMPOSITIONS CONTAINING SAID COMPOUND AND METHODS OF USE



(I)

(57) Abstract

A compound of formula (I) is disclosed which is useful for the treatment of fungal infections and for the treatment of infections caused by *Pneumocystis carinii*.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

### TITLE OF THE INVENTION

CYCLOHEXAPEPTIDYL BISAMINE COMPOUND,  
COMPOSITIONS CONTAINING SAID COMPOUND AND  
METHODS OF USE

### BACKGROUND OF THE INVENTION

The present invention relates to an antifungal and anti-pneumocystis agent of the pneumocandin class, in which the aryl side chain on the 5-aminoethyloxy-4-hydroxyornithine subunit of the hexapeptide consists of a naphthoyl group with an OC<sub>7</sub>H<sub>15</sub> substituent.

Pending US Patent Application Serial No. 07/936,558 filed September 3, 1992 generally discloses pneumocandins containing an aminoethyl ether at the C-5 ornithine position as well as an amine group at the 5th position of the hexapeptide, as having antifungal activity. EP Publication No. 0 535 967 published April 7, 1993 also discloses these pneumocandins.


The compound of the present invention is characterized by an antibiotic spectrum which is largely focused on *Candida*, *Aspergillus*, and *Pneumocystis carinii*. Also, the present compound has safety and pharmacokinetic properties which are superior to other compounds in the class.

There is an increasing need for agents which are effective against opportunistic mycotic infections by such agents as *Candida*, *Aspergillus*, *Cryptococcus* and *Pneumocystis carinii*. The present treatments, i.e., amphotericin B and fluconazole, cause severe side effects or are only fungistatic. The compound of the present invention is considered to be both safe and fungicidal.

- 2 -

SUMMARY OF THE INVENTION

The present invention relates to a novel cyclohexapeptidyl bisamine compound of the formula:



or its acid addition salt.

The present invention also relates to a pharmaceutical composition comprised of the compound in combination with a pharmaceutically acceptable carrier.

Additionally, the present invention relates to a method for treating fungal infections, in particular those caused by *Candida*, *Aspergillus* and *Pneumocystis carinii* which comprises administering the compound of formula I to a patient in need of such treatment in an amount effective to treat the fungal infection. The invention additionally relates to a method for preventing *Pneumocystis carinii* infections in a patient which comprises administration of a preventative amount of the compound of formula I.

- 3 -

Pharmaceutically acceptable salts suitable as acid addition salts as well as salts providing the anion of the quaternary salt are those from acids such as hydrochloric, hydrobromic, phosphoric, sulfuric, maleic, citric, acetic, tartaric, succinic, oxalic, malic, glutamic, pantoic and the like, and include other acids related to the pharmaceutically acceptable salts listed in Journal of Pharmaceutical Science, 66, 2 (1977).

When the compound is a free amine, it is soluble in lower alcohols and polar aprotic solvents such as dimethylformamide (DMF) and pyridine. It is insoluble in solvents such as ether and acetonitrile. When the compound is a quaternary ammonium salt or protonated amine, it is soluble in water and polar solvents.

The compound of the present invention is useful as an antibiotic, especially as an antifungal agent or as an antiparasitic agent. As an antifungal agent, it is useful for the control of both filamentous fungi and yeast. It is especially adaptable to be employed for the treatment of mycotic infections in mammals, especially those caused by *Candida species* such as *C. albicans*, *C. tropicalis*, *C. krusei*, *C. glabrata* and *C. pseudotropicalis*, and *Aspergillus species* such as *A. fumigatus*, *A. flavus* and *A. niger*. It is also useful for the treatment and/or prevention of *Pneumocystis carinii* pneumonia to which immune compromised patients are especially susceptible as hereinafter described.

The previously noted solubility properties are advantageous for utilization in therapeutic applications, especially in injectible compositions.


The compound of the present invention may be obtained from natural products or derivatives of natural products through a sequence of reactions seen in the accompanying reaction schemes or from one of the intermediates which are claimed in concurrently filed copending applications.

The compound of the present invention may be prepared by two different methods. In the first method, illustrated in Scheme 1, the starting material which is represented by formula (E), is a natural product disclosed in US Patent No. 5,202,309 issued April 13, 1993.


- 4 -

The 3-hydroxyglutamine residue may be reduced to a 3-hydroxyornithine residue (Compound F) using any suitable reducing agent. However, preferred agents include a borane complex such as borane with tetrahydrofuran or dimethylsulfide or  $\text{BH}_2\text{Cl}$  with dimethylsulfide or a metal boride such as  $\text{ZrCl}_4/\text{NaBH}_4$ . Compound F has been described in copending application serial number 07/936,561 filed September 3, 1992. Etherification at the 5-position of the 4,5-dihydroxyornithine residue of Compound F may be carried out with ethanolamine and a suitable acid catalyst to give bis-amine Compound G. Compound G has been described in copending application serial number 07/936,558 filed September 3, 1992. The dimethylmyristoyl side chain of Compound G may be deacylated enzymatically with *Comamonas acidovorans* to give triamine Compound H. Compound H is the subject of copending application serial number 08/295,176 filed August 23, 1994. Other suitable enzymatic preparations from a variety of organisms may also be employed for the deacylation. Selective protection of two of the three amine groups of Compound H may be accomplished by any number of reagents. Benzyl 4-nitrophenylcarbonate illustrates this protection step to give Compound J and is shown in Scheme 1. Treatment of Compound J with an activated acylating form of 6-n-heptyloxy-2-naphthoic acid gives the reacylated cyclopeptide K. The preferred activated form is the pentafluorophenyl naphthoate ester. The protecting groups are removed by a suitable method, that being hydrogenolysis for CBz, to give the compound of the present invention.

- 5 -

SCHEME 1


- 6 -



Alternatively, as shown in Scheme 2, the compound of the present invention may be obtained by subjecting Compound E to enzymatic reaction with a preparation of *Comamonas acidovorans* or any suitable organism to give the deacylated cyclopeptide Compound L. Compound L is the subject of US Patent 5,310,873 issued May 10, 1994. Acylation of the free amine group of Compound L may be accomplished using an activated acylating form of 6-n-heptyloxy-2-naphthoic acid. The pentafluorophenyl ester of 6-n-heptyloxy-2-naphthoic acid illustrates the transformation of Compound L to Compound M and is preferred. Reduction of Compound M with a suitable reducing agent such as borane-dimethyl sulfide complex affords amine Compound N. Etherification of Compound N with ethanolamine employing an acid catalyst affords the Compound of the present invention.



- 7 -

SCHEME 2

The compounds of the present invention are active against many fungi and particularly against *Candida*, *Aspergillus* and *Cryptococcus* species. The antifungal properties may be illustrated with

- 8 -

the minimum fungicidal concentration (MFC) determination against certain *Candida* and *Cryptococcus* organisms in a microbroth dilution assay carried out in a Yeast Nitrogen Base (Difco) medium with 1 percent dextrose (YNBD).

In a representative assay, Compound I was solubilized in 100 percent dimethyl sulfoxide (DMSO) at an initial concentration of 5 mg/ml. Once dissolved, the drug stock was brought to a concentration of 512 µg/ml by dilution in water such that the final DMSO concentration was about 10 percent. The solution was then dispensed via a multichannel pipetter into the first column of a 96-well plate (each well containing 0.075 ml of YNBD), resulting in a drug concentration of 256 µg/ml. Compounds in the first column were diluted 2-fold across the rows yielding final drug concentrations ranging from 256 µg/ml to 0.12 µg/ml.

Four-hour broth cultures of organisms to be tested were adjusted using a spectrophotometer at 600 nm to equal a 0.5 McFarland Standard. This suspension was diluted 1:100 in YNBD to yield a cell concentration of  $1.5 \times 10^4$  colony forming units (CFU)/ml. Aliquots of the suspension (0.075 ml) were inoculated into each well of the microtiter plate resulting in a final cell inoculum of  $5.25 \times 10^3$  CFU/ml and final drug concentrations ranging from 128 µg/ml to 0.06 µg/ml. Each assay includes one row for drug-free control wells and one row for cell-free control wells.

After 24 hours of incubation, the microtiter plates were shaken gently on a shaker to resuspend the cells. The MIC-2000 inoculator was used to transfer a 1.5 microliter sample from each well of the 96-well microtiter plate to a single reservoir inoculum plate containing Sabouraud dextrose agar (SDA). The inoculated SDA plates were incubated for 24 hours at 35°C. However, for *Cryptococcus neoformans* strains, SDA plates were inoculated at 48 hours and incubated 48 hours after being spotted on SDA before making minimum fungicidal concentration (MFC) readings. The results were as follows:

- 9 -

Organism		MFC μg/mL
<i>C. albicans</i>	CLY 539	<0.06
<i>C. albicans</i>	MY 1055	<0.06
<i>C. glabrata</i>	MY 1381	<0.06
<i>C. guilliermondii</i>	MY 1019	0.25
<i>C. krusei</i>		<0.06
<i>C. parapsilosis</i>	MY 1010	<0.06
<i>C. pseudotropicalis</i>	MY 2099	<0.06
<i>C. tropicalis</i>	MY 1012	<0.06
<i>Cr. neoformans</i>	MY 2061	128
<i>Cr. neoformans</i>	MY 2062	32

Compound I also shows *in vivo* effectiveness against fungi.

Growth from an overnight SDA culture of *Candida albicans* MY 1055 was suspended in sterile saline and the cell concentration determined by hemacytometer count and the cell suspension adjusted to  $3.75 \times 10^5$  cells/ml. Then 0.2 milliliter of this suspension was administered I.V. in the tail vein of mice so that the final inoculum was  $7.5 \times 10^4$  cells/mouse.

The assay then was carried out by administering aqueous solutions of Compound I at various concentrations intraperitoneally (I.P.), twice daily (b.i.d.) for two consecutive days to 18 to 20 gram female DBA/2 mice, which previously had been infected with *Candida albicans* in the manner described above. Distilled water was administered I.P. to *C. albicans* challenged mice as controls. After four days, the mice were sacrificed by carbon dioxide gas, paired kidneys were removed aseptically and placed in sterile polyethylene bags containing 5 milliliters of sterile saline. The kidneys were homogenized in the bags, serially diluted in sterile saline and aliquots spread on the surface of SDA plates. The plates were incubated at 35°C for 48 hours and yeast colonies were enumerated for determination of colony forming units (CFU) per gram of kidneys. Compound I showed greater

- 10 -

than 99 percent reduction of recoverable *Candida* CFUs when dosed at 0.375 and 0.09 mg/kg I.P. as described above.

The compound of the present invention may also be useful for inhibiting or alleviating *Pneumocystis carinii* infections in immune compromised patients. The efficacy of the compound of the present invention for therapeutic or anti-infective purposes may be demonstrated in studies on immunosuppressed mice.

In a representative study, the effectiveness of Compound I was determined. Mice (weighing approximately 20 grams) were immunosuppressed with dexamethasone in the drinking water (2.0 mg/L) and maintained on a low protein diet for seven weeks to induce the development of *Pneumocystis* pneumonia from a latent infection. Before drug treatment, two mice were sacrificed to confirm the presence of *Pneumocystis carinii* pneumonia (PCP); both mice were found to have infections. Five mice (weighing approximately 20 grams) were injected twice daily for four days subcutaneously (sc) with the compound in 0.25 ml of vehicle (distilled water). A vehicle control was also carried out. All animals continued to receive dexamethasone in the drinking water and low protein diet during the treatment period. At the completion of the treatment, all animals were sacrificed, the lungs were removed and processed, and the extent of disease determined by microscopic analysis of stained slides. The results of this study showed Compound I was >95 percent effective in reducing *P. carinii* cysts in 5 mice when dosed at 0.125 mg/kg.

The compound of the present invention has surprising and unexpected safety benefits in that there is markedly improved acute tolerability in mammals than with other related compounds. To demonstrate, laboratory test animals (rhesus monkeys) were dosed with the compound intravenously and observed for any abnormal effects. Unexpectedly, no notable response was observed for the compound of the present invention when dosed at identical levels of related compounds that do cause cardiovascular effects.

- 11 -

Additionally, the acute tolerability of the compound of the present invention in mice has been shown to be superior to that of related compounds.

The outstanding properties are most effectively utilized when Compound I is formulated into novel pharmaceutical compositions with a pharmaceutically acceptable carrier according to conventional pharmaceutical compounding techniques.

The novel compositions contain at least a therapeutic antifungal or antipneumocystis amount of the active compound of the present invention. Generally, the composition contains at least 1 percent by weight of Compound I. Concentrate compositions suitable for dilutions prior to use may contain 90 percent or more by weight. The compositions include compositions suitable for oral, topical, parenteral (including intraperitoneal, subcutaneous, intramuscular, and intravenous), nasal, and suppository administration, or insufflation. The compositions may be prepacked by intimately mixing Compound I with the components suitable for the medium desired.

Compositions formulated for oral administration may be a liquid composition or a solid composition. For liquid preparations, the therapeutic agent may be formulated with liquid carriers such as water, glycols, oils, alcohols, and the like, and for solid preparations such as capsules and tablets, with solid carriers such as starches, sugars, kaolin, ethyl cellulose, calcium and sodium carbonate, calcium phosphate, kaolin, talc, lactose, generally with a lubricant such as calcium stearate, together with binders disintegrating agents and the like. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage form. It is especially advantageous to formulate the compositions in unit dosage form (as hereinafter defined) for ease of administration and uniformity of dosage. Compositions in unit dosage form constitute an aspect of the present invention.

Compositions may be formulated for injection and for injection take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles such as 0.85 percent sodium chloride or 5 percent dextrose in water and may contain formulating agents such as

- 12 -

suspending, stabilizing and/or dispersing agents. Buffering agents as well as additives such as saline or glucose may be added to make the solutions isotonic. Compound I may also be solubilized in alcohol/propylene glycol or polyethylene glycol for drip intravenous administration. These compositions also may be presented in unit dosage form in ampoules or in multidose containers, preferably with added preservative. Alternatively, the active ingredients may be in powder form for reconstituting with a suitable vehicle prior to administration.

The term "unit dosage form" as used in the specification and claims refer to physically discrete units, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the pharmaceutical carrier. Examples of such unit dosage forms are tablets, capsules, pills, powder packets, wafers, measured units in ampoules or in multidose containers and the like. A unit dosage of the present invention will generally contain from 10 to 500 milligrams of the compound.

When the compound of the present invention is for antifungal use, any method of administration may be employed.

When Compound I is to be employed for control of pneumocystis infections any method may be employed although it may be desirable to directly treat lung and bronchi. In such administration inhalation methods are employed. For administration by inhalation, the compounds of the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or nebulisers. The preferred delivery system for inhalation is a metered dose inhalation (MDI) aerosol, which may be formulated as a suspension or solution of the compound in suitable propellants, such as fluorocarbons or hydrocarbons.

Although the compound of the present invention may be employed as tablets, capsules, topical compositions, insufflation powders, suppositories and the like, the solubility of the compounds of the present invention in water and aqueous media render them adaptable

- 13 -

for use in injectible formulations and also in liquid compositions suitable for aerosol sprays.

The invention is further described in connection with the following non-limiting examples.

### EXAMPLE 1

#### Part A. Preparation of Compound F

Compound E (15.9 g, 89 area% pure, 3.4 wt% water, 0.0128 mol) was added to dry THF (0.64 L) and the suspension was dried to <10 mol% water by refluxing through a bed of 3Å molecular sieves. Additional dry THF was added to reconstitute the mixture to the original volume and the suspension was cooled to < 4°C with an ice/water/methanol bath.

Neat BH<sub>3</sub>•SMe<sub>2</sub> (10.91 g, 0.144 mol) was added over ten minutes and the reaction mixture was maintained at 0-4°C. The reaction progress was monitored by HPLC until the ratio of starting material to product was 1:1 indicating the end of the reaction (3.5 h). At 4 hours, the mixture was cooled to -12°C and slowly quenched with 2N HCl (0.036 L). This solution was diluted to 1.14 L with water. The assay yield of Compound F was 6.60 g (47%).

The quenched solution was diluted to 4 L and loaded onto a medium-pressure column of LiChroprep RP-C18 adsorbent (158 g). After loading, the column was washed with 1.2 L water and the amine was eluted with 1.9 L of 1:4 v/v acetonitrile/water, and then 0.38 L of 1:3 v/v acetonitrile/water.

The rich cuts (> 80 area%) were combined and diluted with water to a 1:7.3 v/v acetonitrile/water solution (1.70 L total). This mixture was loaded to the same column described above, and the column was washed with 0.57 L water. The desired compound was eluted with 0.57 L methanol. The rich cut fractions (> 85 area%) were combined and concentrated by rotary evaporation and static high vacuum to give 6.81 g (87 wt% pure, 6.8 wt% water) containing 5.92 g

- 14 -

of Compound F (where R<sup>1</sup> is dimethyltridecyl) hydrochloride salt for an isolated yield of 43%.

<sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD): δ 7.12 (d, 2H), 6.75 (d, 2H), 5.18 (d, 1H), 4.97 (d, 1H), 1.19 (d, 3H), 0.89 (t, 3H), 0.86 (d, 6H).

Mass spectrum (FAB) *m/z* (M+Li)<sup>+</sup>: 1058.

#### Part B. Preparation of Compound G

A solution of Compound F as the hydrochloride salt (41.4g, 0.038 mol), ethanolamine hydrochloride (100 g, 1.0 mol), and 4.0M hydrochloric acid in 1,4-dioxane (11.4 ml, 0.046 mol) in anhydrous dimethylsulfoxide (200 ml) was stirred at ambient temperature under a nitrogen atmosphere for a period of 66 hours. HPLC analysis (41% acetonitrile/water/0.1% trifluoroacetic acid) showed 6% unreacted Compound F (*t<sub>R</sub>* = 7.9 min) and 56% Compound G (*t<sub>R</sub>* = 4.2 min). The reaction mixture was diluted to 1 liter with water and filtered through a bed of filter aid. Pump-injection of the filtrate onto a preparative C18 HPLC column was followed by elution with 5-15% acetonitrile/water/0.1% acetic acid in 5% step gradients. The product-containing fractions (40% acetonitrile/water/0.1% trifluoroacetic acid, *t<sub>R</sub>* = 4.2 min) were combined and diluted 10% with water. Compound G was isolated by C18 solid-phase extraction eluting with methanol. Removal of the methanol *in vacuo* provided a syrup which was dissolved in 50% CH<sub>3</sub>CN/water and lyophilized to give 18.1 g of Compound G as a mixture of hydrochloride and acetate salts suitable for use in the subsequent deacylation step. HPLC purity > 80%. Rechromatography of a portion of the mixed salt followed by anion exchange to the dihydrochloride provided Compound G as the dihydrochloride salt. HPLC purity > 99%.

<sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD): δ 7.12 (d, 2H), 6.77 (d, 2H), 5.18 (d, 1H), 3.14 (t, 2H), 3.09 (t, 2H).



- 15 -

Anal. (C<sub>52</sub>H<sub>89</sub>Cl<sub>2</sub>N<sub>9</sub>O<sub>16</sub>) C, H, N, Cl. HRMS (FAB) *m/z* (M+1)  
Calculated for C<sub>52</sub>H<sub>88</sub>N<sub>9</sub>O<sub>16</sub> 1094.6348  
Found 1094.6338.

### Part C. Deacylation of Compound G to Give Triamine H

Compound G was enzymatically deacylated as follows:

#### i. Preparation of *C. acidovorans* cells

To establish a seed culture of *C. acidovorans* ATCC 53942, a loopful of cells from an LB agar slant of the organism was used to inoculate 50 mL of Lauria-Bertani (LB) medium in a 250-mL unbaffled Erlenmeyer flask. The flask was incubated at 25°C with agitation at 220 rpm for about 18 hours and 0.1 mL of this culture was used to inoculate each of three 250-mL Erlenmeyer flasks containing 50 mL of LB medium. After incubating under the same conditions for 18 hours, 20 mL of the grown cells were used to inoculate 15 liters of LB medium in each of six 23-liter fermentors. The fermentors were operated at 30°C with agitation at 400 rpm and aeration at 7.4 L/min for 18 hours. Cultures from three fermentors were combined, washed and concentrated to 11 liters in 50 mM potassium phosphate buffer, pH 7.5 using a hollow fiber ultrafiltration system.

#### ii. Deacylation of Compound G

Deacylation was set up in two 15-liter Microferm fermentors (New Brunswick Scientific). In one vessel (A), 10 g of Compound G dissolved in 250 mL of distilled water was added to 5 liters of the washed and buffered *C. acidovorans* cells while in the second vessel (B), 8.2 g of Compound G dissolved in 200 mL of distilled water was added to 4 liters of the cells. Addition of Compound G to each vessel was accomplished with a pipette over a 5-minute period. To reduce foaming, about 2 mL and 1.5 mL of P2000 were added to vessels A and B respectively. Each fermentor was operated with stirring at 100 rpm, aeration at 1.5 L/min and temperature at 37°C for the first 2.5 hours and 30°C for the last 21.5 hours. After 24 hours,

- 16 -

HPLC analyses indicated that deacylation was 100% complete in each vessel. The contents of both vessels were combined and the cells were removed by hollow fiber filtration. HPLC analysis confirmed that the filtrate contained Compound H, the nucleus of Compound G.

### iii. Isolation of Compound H

The deacylation broth from the procedure above was processed by loading 17 L broth onto 4x120 g resin columns (BioRad AG50W-X8, H<sup>+</sup> form). The columns were each washed with 1 L of water to pH = 7.0 prior to loading the phosphate-buffered broth. Each loading was followed by 1 L water wash and compound H was then eluted with 0.15N NH<sub>4</sub>OH. The product-containing fractions (approximately 2L) were combined and lyophilized to give 10.3 g of crude Compound H (free amine). This was then slurried in 100 mL of water, cooled to 0 °C and 100 mL of 2% CF<sub>3</sub>COOH/H<sub>2</sub>O was added (5 eq, pH = 2). This was then loaded onto a 50 g RP-C18 plug (rinsing with 0.1% aqueous trifluoroacetic acid). Lyophilization then gave 16.2 g of Compound H as the tri-TFA salt (~70% pure). A small portion was further purified on an Rx-C18 column (21.2 mm x 25 cm) using 5% CH<sub>3</sub>CN/H<sub>2</sub>O (0.5% CF<sub>3</sub>COOH). Elution at a flow rate of 10 ml/min was monitored by UV at 220 and 277 nm. The product-containing fractions, as determined by analytical C18-HPLC (4.6 mm x 25 cm, 3.5% CH<sub>3</sub>CN/H<sub>2</sub>O (0.1% CF<sub>3</sub>COOH) at 1.5 ml/min, t<sub>R</sub> = 4.9 min) were combined and lyophilized to give >98% pure Compound H.

<sup>1</sup>H NMR (500MHz,CD<sub>3</sub>OD) : δ 7.13 (d, J=8.4 Hz, 2H), 6.75 (d, J=8.7 Hz, 2H), 5.24 (d, J=3.2 Hz, 1H), 5.01 (d, J=2.7 Hz, 1H), 4.42 (dd, J=5.5 and 11.2 Hz, 1H), 4.38 (d, J=7.8 Hz, 1H), 3.98 (dd, 3.1 and 11.1 Hz, 1H), 3.15 (m, 2H), 3.05 (t, J=7.3 Hz, 2H), 2.43(dd, J=6.9 and 13.0 Hz, 1H), 1.26 (d, J=6.2 Hz, 3H).

ESI-MS (M+H)<sup>+</sup> = 857.6

- 17 -

iv. Preparation of Compound I from Compound HStep a.

To a stirred solution of Compound H (16.0 g [ $\sim 70\%$  pure], 0.0094 mol) in anhydrous dimethylformamide (75 mL) was added triethylamine (9.3 mL, 0.067 mol), followed by benzyl-4-nitrophenyl carbonate (5.11 g, 0.019 mol), and this was stirred for 20 minutes. HPLC indicated incomplete formation of Compound J (analytical C18-HPLC : 30% acetonitrile/water/0.1% trifluoroacetic acid,  $t_R = 3.7$  min). Additional carbonate (1.46 g, 0.005 mol) was added and HPLC indicated mainly Compound J as product. The pentafluorophenyl 6-n-heptyloxy-2-napthoate (6.65 g, 0.015 mol) was then added and stirred for 18 hours. HPLC indicated consumption of Compound J and formation of acylated product, Compound K. The reaction was diluted with water (75 mL) and loaded onto a RP C18 flash column (50 g) packed in 30%  $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ ; rinsed with 6 x 50 mL of 50%  $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ ; and lastly 10 x 50 mL 100% methanol. The product-containing methanol fractions were combined and concentrated to give 11.1 g of crude Compound K (80% pure). A small portion of Compound K was further purified on Zorbax Rx-C18 (21.2 x 250 mm) eluting with 70%  $\text{CH}_3\text{CN}/\text{H}_2\text{O}$  at a flow rate of 10 mL/min. The product-containing fractions, as determined by HPLC (75% A/B,  $t_R = 3.1$  min), were combined and lyophilized to give  $> 98\%$  pure Compound K.

$^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ) :  $\delta$  8.32 (s, 1H), 7.85 (d,  $J=7.6$  Hz, 1H), 7.74 (d,  $J=8.9$  Hz, 1H), 7.66 (d,  $J=8.7$  Hz, 1H), 7.10-7.30 (m, 14H), 6.75 (d,  $J=8.5$  Hz, 2H), 5.29 (s, 1H), 4.99 (m, 2H), 4.93 (s, 4H), 3.96 (dd,  $J=3.1$  and 11.1 Hz, 1H), 3.61 (m, 1H), 3.52 (m, 1H), 3.40 (m, 1H), 2.43 (dd,  $J=6.9$  and 13.0 Hz, 1H), 1.84 (m, 2H), 1.22 (d,  $J=6.1$  Hz, 3H), 0.92 (t,  $J=6.9$  Hz, 3H).

ESI-MS ( $\text{M}+\text{TFA}$ ) $^-$  = 1505

- 18 -

Step b.

To a stirred suspension of Compound K (11.0 g [80% pure], 0.006 mol) in CH<sub>3</sub>OH (200 mL) and CH<sub>3</sub>COOH (20 mL) was added 5.5g 10% Pd/C. This was then subjected to an atmosphere of hydrogen (balloon pressure) and allowed to stir for 3 hours. HPLC indicated complete deprotection and catalyst was filtered through a pad of diatomaceous earth, rinsing with CH<sub>3</sub>OH, and concentration gave crude Compound I. This was then dissolved in water (150 mL) and loaded onto Delta Prep C18 for purification. A step-gradient chromatography from 10-30% acetonitrile/water/0.1% acetic acid was used and lyophilization of the appropriate fractions as determined by HPLC (35% acetonitrile/water/0.1% trifluoroacetic acid,  $t_R$  = 4.7 min) gave 6.2 g of Compound I as the di-acetate salt. This was then dissolved in water (200 mL) and applied to a BioRad AG2-X8, Cl<sup>-</sup> resin column (70 g); elution with H<sub>2</sub>O and subsequent lyophilization gave 5.8 g of Compound I as the dihydrochloride salt.

<sup>1</sup>H NMR (CD<sub>3</sub>OD) :  $\delta$  8.36 (s, 1H), 7.80-7.90 (m, 3H), 7.28 (d, J=1.8 Hz, 1H), 7.21 (dd, J=2.2 and 9.0 Hz, 1H), 7.13 (d, J=8.5 Hz, 2H), 6.76 (d, J=8.5 Hz, 2H), 5.29 (s(br), 1H), 5.03 (d, J=3.0 Hz, 1H), 4.94 (d, J=6.6 Hz, 2H), 4.76 (dd, J=4.7 and 12.9 Hz, 1H), 4.13 (m, 3H), 3.87 (t, J=6.8 Hz, 2H), 3.18 (m, 2H), 3.10 (t, J=5.5 Hz, 2H), 2.45 (dd, J=7.0 and 12.7 Hz, 1H), 1.53 (m, 2H), 1.22 (d, J=6.2 Hz, 3H), 0.92 (t, J=6.9 Hz, 3H).

Anal. (C<sub>54</sub>H<sub>79</sub>Cl<sub>2</sub>N<sub>9</sub>O<sub>17</sub>) C, H, N, Cl. ESI-MS (M+H)<sup>+</sup> = 1094.6

EXAMPLE 2Part A. Deacylation of Pneumocandin B<sub>0</sub>

The preparation of the deacylating enzyme is first described. A loopful of *Comamonas acidovorans* ATCC 53942 is inoculated into 50 milliliters of Luria Bertani medium of the following

- 19 -

composition: per liter Bacto-Trypton, 10 g; Bacto-Yeast Extract, 5 g and sodium chloride 10 g and solidified with 2 percent agar and incubating for 24 hours with shaking to obtain a seed culture. Cells for the deacylation are then grown by diluting a 50 milliliter portion of seed culture 1:500 into fresh Luria-Bertani medium and incubating at 25 °C with shaking for 16 hours. Cells are then harvested by centrifugation, washed by resuspending in 1 percent sodium chloride and centrifuged and then resuspended in potassium phosphate buffer at pH 6.5.

A solution of pneumocandin B<sub>0</sub> (Compound E) in dimethylsulfoxide is then added to a stirred suspension of the cells of *C. acidovorans* thus obtained and the mixture maintained at 37 °C for 18 hours to obtain the deacylated product, Compound L, which is recovered in the supernatant after centrifugation.

Compound L may be isolated by adsorbing the supernatant on HP-20 resin with water, eluting with methanol and concentrating the eluates. The eluates are combined, diluted with water, charged to a preparative HPLC system equipped with a Whatman Partisil 10 SCX column and then eluted with 0.01M potassium phosphate (pH=6) buffer and monitored via UV at 210 nm. Cuts rich in the deacylated product were combined and the resulting mixture absorbed and eluted from HP-20 resin with methanol to remove buffer salts and to obtain Compound L with a molecular weight of 826.

#### Part B. Reacylation of Compound L to Give Compound M

Pentafluorophenyl 6-n-heptyloxy-2-naphthoate is prepared as follows. To a suspension of 6-n-heptyloxy-2-naphthoic acid (10.5 mmol) and dicyclohexylcarbodiimide (10.5 mmol) in ethyl acetate at 0°C is added pentafluorophenol (11.5 mmol). The mixture is stirred at 25 °C for a period of 18 h. Any precipitate is removed by filtration. The filtrate is washed twice with water and brine and dried with magnesium sulfate. Removal of the ethyl acetate *in vacuo* gives the desired pentafluorophenyl 6-n-heptyloxy-2-naphthoate.

- 20 -

Pentafluorophenyl 6-n-heptyloxy-2-naphthoate (1.1 mmol) prepared as above is added to a solution of Compound L from Part A (1.0 mmol) in anhydrous N,N-dimethylformamide. The resulting solution is stirred at room temperature for a period of 12 to 24 h. The mixture is subsequently diluted with water. C18-Flash chromatography of the mixture eluting with 30-100% CH<sub>3</sub>CN/H<sub>2</sub>O in 10% step gradients followed by lyophilization of the product-containing fractions as determined by analytical HPLC (Zorbax RX-C18, UV detection at 210 nm) gives the desired Compound M with a molecular weight of 1095.2

#### Part C. Reduction of Compound M to Give Compound N

A reaction vessel is outfitted with an addition funnel charged with 3-4 angstrom molecular sieves and a reflux condenser above the addition funnel. Within the reaction vessel, Compound M is suspended in dry tetrahydrofuran and heated to reflux under a nitrogen atmosphere. The mixture is refluxed for 12-24 hours, then cooled to 0°C. Borane-methylsulfide complex (10-20 equivalents) is added dropwise over 1-2 hours. The mixture is stirred at 0°C for 2-5 hours and then carefully quenched with 2N hydrochloric acid. Stirring is continued for 1 hour after the final addition of hydrochloric acid. The mixture is concentrated to 40-50 percent of the original volume and then diluted with 6-7 volumes of distilled water. Purification by preparative reverse phase HPLC (DeltaPak C18, acetonitrile/water/0.1% acetic acid, UV detection at 210 nm) and subsequent lyophilization of the appropriate fractions, as determined by analytical HPLC, gives the desired compound. The hydrochloride salt is prepared by passage of an aqueous solution of the residue through an anion exchange column (BioRad AG2-X8, Cl<sup>-</sup> resin) to give the desired Compound N with a molecular weight of 1117.6.

- 21 -

Part D. Aminoethyletherification of Compound N to Compound I

To a solution of the cyclohexapeptidyl propanolamine compound (N) (0.2 mmol) in DMSO (8 mL) containing 2-aminoethanol as the hydrochloride salt (20 mmol, 100 eq based on the cyclohexapeptide propanolamine) is added 4M HCl in dioxane (0.4 mmol) and the mixture is stirred at room temperature for two to seven days. The progress of the reaction may be monitored by HPLC using a 0.5 cm X 25 cm analytical Zorbax C18 column eluted with a mixture of acetonitrile and water containing 0.1 % trifluoroacetic acid (typically the solvent was from 10-50% acetonitrile). When the reaction is judged to be complete, the reaction mixture is diluted with five to ten volumes of water and the resulting mixture applied to a reverse phase flash silica gel column using Lichroprep C18 (particle size 40-63 microns). The mixture is then eluted initially with 5%-15% acetonitrile in water to remove the excess DMSO and 2-aminoethanol hydrochloride, then with 10-50% acetonitrile in water to elute the product. Fractions containing the desired product are combined, concentrated and lyophilized to afford the product. Subsequent exchange for chloride ion may be accomplished by dissolving the residue in water and passing through a BioRad AG2-X8, Cl<sup>-</sup> resin column to give the desired Compound I as the dihydrochloride with a molecular weight of 1197.2.





- 23 -

5. A pharmaceutical composition as claimed in Claim 4 where the acid addition salt is selected from the group consisting of acetate, trifluoroacetate, chloride, bromide, tartrate, succinate, glutamate, sulfate, phosphate and pamoate.

6. A pharmaceutical composition as claimed in Claim 5 wherein the acid addition salt is trifluoroacetate or hydrochloride.

7. A method for treating fungal infections in a patient in need of such treatment, comprising administering to said patient an effective amount of the compound as defined in Claim 1.

8. A method for treating fungal infections in a patient in need of such treatment, comprising administering to said patient an effective amount of a compound as defined in Claim 2.

9. A method for treating fungal infections in a patient in need of such treatment, comprising administering to said patient an effective amount of a compound as defined in Claim 3.

10. A method for treating an infection caused by *Candida* sp. in a patient which comprises administering to said patient an effective amount of a compound as defined in Claim 1.

11. A method for treating an infection caused by *Candida* sp. in a patient which comprises administering to said patient an effective amount of a compound as defined in Claim 2.

12. A method for treating an infection caused by *Candida* sp. in a patient which comprises administering to said patient an effective amount of a compound as defined in Claim 3.

- 24 -

13. A method for treating an infection caused by *Aspergillus sp.* in a patient which comprises administering to said patient an effective amount of a compound as defined in Claim 1.

14. A method for treating an infection caused by *Aspergillus sp.* in a patient which comprises administering to said patient an effective amount of a compound as defined in Claim 2.

15. A method for treating an infection caused by *Aspergillus sp.* in a patient which comprises administering to said patient an effective amount of a compound as defined in Claim 3.

16. A method for treating or preventing an infection or condition caused by *Pneumocystis carinii* in a patient in need of such treatment or prevention which comprises administering to said patient a preventative or therapeutic amount of the compound of Claim 1.

17. A method for treating or preventing an infection or condition caused by *Pneumocystis carinii* in a patient in need of such treatment or prevention which comprises administering to said patient a preventative or therapeutic amount of the compound of Claim 2.

18. A method for treating or preventing an infection or condition caused by *Pneumocystis carinii* in a patient in need of such treatment or prevention which comprises administering to said patient a preventative or therapeutic amount of the compound of Claim 3.

# INTERNATIONAL SEARCH REPORT

International Application No  
PC1/US 96/17715

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C07K/56 A61K38/12

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 535 967 A (MERCK & CO INC) 7 April 1993 cited in the application see page 12; claims 1-7; example VIIA ---	1-18
Y	EP 0 644 199 A (FUJISAWA PHARMACEUTICAL CO) 22 March 1995 see examples 3,11,14,20,21 ---	1-18
A	WO 94 21677 A (MERCK & CO INC ;BALKOVEC JAMES M (US); BOUFFARD FRANCES AILEEN (US) 29 September 1994 See especially example 16 and the table on page 19 -----	1-18

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- \*A\* document member of the same patent family

Date of the actual completion of the international search

20 February 1997

Date of mailing of the international search report

10.03.97

Name and mailing address of the ISA  
European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+ 31-70) 340-3016

Authorized officer

Groenendijk, M

# INTERNATIONAL SEARCH REPORT

national application No.

PCT/US 96/17715

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 7-18  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claim(s) 7-18  
is(are) directed to a method of treatment of the human/animal  
body, the search has been carried out and based on the alleged  
effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 96/17715

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
EP-A-0535967	07-04-93	CA-A-	2079187	02-04-93
		JP-A-	5239094	17-09-93
		JP-B-	7094470	11-10-95
-----				
EP-A-0644199	22-03-95	AU-A-	6199494	24-11-94
		CA-A-	2123921	18-11-94
		CN-A-	1100104	15-03-95
		HU-A-	68385	28-06-95
		JP-A-	6340693	13-12-94
		US-A-	5569646	29-10-96
		ZA-A-	9403356	28-03-95
-----				
WO-A-9421677	29-09-94	US-A-	5378804	03-01-95
		AU-B-	668804	16-05-96
		AU-A-	5783294	22-09-94
		BG-A-	99999	30-04-96
		BR-A-	9406009	26-12-95
		CA-A-	2118757	17-09-94
		CN-A-	1119441	27-03-96
		CZ-A-	9502358	14-02-96
		EP-A-	0620232	19-10-94
		FI-A-	954327	14-09-95
		HR-A-	940161	31-08-96
		HU-A-	9500565	30-10-95
		HU-A-	73496	28-08-96
		JP-A-	6321986	22-11-94
		NO-A-	953648	15-11-95
		PL-A-	310634	27-12-95
		SI-A-	9420013	31-08-96
		SK-A-	113695	07-02-96
		US-A-	5514650	07-05-96
		ZA-A-	9401807	13-10-94
-----				